

Equivalent martingale measures in Lévy markets

João Guerra

CEMAPRE and ISEG, UTL

November, 12, 2014

From the previous lecture

- If the process $e^Y = (e^{Y(t)}, t \geq 0)$ is a martingale then

$$\begin{aligned} e^{Y(t)} = & 1 + \int_0^t e^{Y(s-)} F(s) dB(s) + \int_0^t \int_{|x| < 1} e^{Y(s-)} \left(e^{H(s,x)} - 1 \right) \tilde{N}(ds, dx) \\ & + \int_0^t \int_{|x| \geq 1} e^{Y(s-)} \left(e^{K(s,x)} - 1 \right) \tilde{N}(ds, dx) \end{aligned} \quad (1)$$

- \tilde{S} is a Q-martingale if and only if

$$m\sigma + \mu - r + k\sigma F(t) + \sigma \int_{\mathbb{R} - \{0\}} x \left(e^{H(t,x)} - 1 \right) \nu(dx) = 0 \text{ a.s.} \quad (2)$$

Incomplete markets and Esscher transform

- Equivalent measures Q exist with respect to which \tilde{S} will be a martingale, but these will no longer be unique in general
- We must follow a selection principle to reduce the class of all possible measures Q to a subclass, within which a unique measure can be found.
- Additional assumption:

$$\int_{|x| \geq 1} e^{ux} \nu(dx) < \infty$$

for all $u \in \mathbb{R}$.

- In this case, we can analytically continue the Lévy- Khintchine formula to obtain

$$\mathbb{E} \left[e^{-uX(t)} \right] = e^{-t\psi(u)}$$

where

$$\psi(u) = -\eta(iu) = bu - \frac{1}{2}k^2u^2 + \int_c^\infty (1 - e^{-ux} - ux\mathbf{1}_{\{|x| < 1\}}(x)) \nu(dx).$$

Incomplete markets and Esscher transform

- The processes

$$M_u(t) = \exp(iuX(t) - t\eta(u)),$$

$$N_u(t) = M_{iu}(t) = \exp(-uX(t) + t\psi(u))$$

are martingales and N_u is strictly positive.

- Define a new probability measure by

$$\frac{dQ_u}{dP} \Big|_{\mathcal{F}_t} = N_u(t).$$

- Q_u is called the Esscher transform of P by N_u .

Incomplete markets and Esscher transform

- Applying the Itô formula to N_u , we have

$$dN_u(t) = N_u(t-) \left(-kuB(t) + (e^{-ux} - 1) \tilde{N}(dt, dx) \right).$$

- Comparing this with (1) for exponential martingales e^Y , we have that

$$\begin{aligned} F(t) &= -ku, \\ H(t, x) &= -ux \end{aligned}$$

and the condition for Q_u to be a martingale (2) is

$$m\sigma + \mu - r - k^2u\sigma + \sigma \int_c^\infty x (e^{-ux} - 1) \nu(dx) = 0 \text{ a.s.}$$

Incomplete markets and Esscher transform

- Let $z(u) = \int_c^\infty x (e^{-ux} - 1) \nu(dx) - k^2u$. Then the martingale condition is:

$$z(u) = \frac{r - \mu - m\sigma}{\sigma}. \quad (3)$$

- Since $z'(u) < 0$, z is strictly decreasing, and therefore there is a unique u (a unique measure Q_u) that satisfies (3).
- The Esscher transform is such that this measure Q_u minimizes the relative entropy $H(Q|P)$ between the measures Q and P (a measure of "distance" between two measures), where

$$H(Q|P) = \mathbb{E}_Q \left[\ln \left(\frac{dQ}{dP} \right) \right] = \mathbb{E}_P \left[\frac{dQ}{dP} \ln \left(\frac{dQ}{dP} \right) \right].$$

Absence of arbitrage in exponential Lévy models

- Let X be a Lévy process and consider a market model where $S_t = S_0 \exp(X_t)$.
- Theorem (see Cont and Tankov, pages 310-311): If the trajectories of X are neither increasing (a.s.) nor decreasing (a.s.), then the exponential Lévy market model given by $S_t = S_0 \exp(X_t)$ is arbitrage free: there exists a probability measure Q equivalent to P (equivalent martingale measure) such that $\tilde{S}_t = e^{-rt} S_t$ is a Q -martingale.
- In other words, the exponential-Lévy model is arbitrage free in the following cases (not mutually exclusive):
 - 1) X has a nonzero Gaussian component (or diffusion coeff.): $\sigma > 0$.
 - 2) X has infinite variation: $\int_{|x| < 1} |x| \nu(dx) = \infty$.
 - 3) X has both positive and negative jumps.
 - 4) X has positive jumps and negative drift or negative jumps and positive drift.

The mean-correcting martingale measure

- A practical way to obtain an equivalent martingale measure Q in a exponential Lévy model of type $S_t = S_0 \exp(X_t)$, is by mean correcting the exponential of a Lévy process (see Schoutens, pages 79-80).
- We can correct the exponential of the Lévy process X , by adding a new drift term mt (with new parameter m):

$$\bar{X}_t = mt + X_t.$$

- When comparing the characteristics triplet of \bar{X} with those of X , the only parameter that changes is the drift: $\bar{b} = b + m$.

The mean-correcting martingale measure

- We can change the m parameter of the process X such that $\tilde{S}_t = e^{-rt} S_t$ is a martingale. This is equivalent to choose an equivalent martingale measure Q .
- Example: in the Black-Scholes model, we change the mean of the normal distribution $\mu - \frac{1}{2}\sigma^2 = m_{old}$ (the m_{old}) into the new m parameter:

$$m_{new} = r - \frac{1}{2}\sigma^2,$$

or

$$m_{new} = m_{old} + r - \ln[\phi(-i)],$$

where $\phi(x)$ is the characteristic function of the log-returns involving the m_{old} parameter.

In the Black-Scholes model, $\ln[\phi(-i)] = \mu$.

- This choice of m_{new} will imply that the discounted price $\tilde{S}_t = e^{-rt} S_t$ is a martingale.

The mean-correcting martingale measure





- Procedure:
 - 1) Estimate in some way the parameters involved in the process.
 - 2) Then change the m parameter in a way that

$$m_{new} = m_{old} + r - \ln[\phi(-i)],$$

where $\phi(x)$ is the characteristic function of the log-returns involving the m_{old} parameter.

3) Then, with this new m_{new} parameter in the Lévy process, the discounted price $\tilde{S}_t = e^{-rt} S_t$ is a martingale and we have chosen the mean-correcting equivalent martingale measure.

- In page 78 of Schoutens, the author lists what is the value of the m parameter for several Lévy processes (CGMY, VG, NIG, etc...)

-  Applebaum, D. (2004). Lévy Processes and Stochastic Calculus. Cambridge University Press. - (Section 5.6)
-  Applebaum, D. (2005). Lectures on Lévy Processes, Stochastic Calculus and Financial Applications, Ovronnaz September 2005, Lecture 3 in <http://www.applebaum.staff.shef.ac.uk/ovron3.pdf>
-  Cont, R. and Tankov, P. (2003). Financial modelling with jump processes. Chapman and Hall/CRC Press, (Section 9.5.)
-  Schoutens, W. (2002). Lévy Processes in Finance: Pricing Financial Derivatives. Wiley, (Sections 5.4 and 6.2.)