

Exercises from  
*Arbitrage Theory in Continuous Time (3:rd ed)*  
Chapters 4-5

Tomas Björk  
Stockholm School of Economics

May 10, 2010

## 1 Exercises

**Exercise 1.1** Compute the stochastic differential  $dZ$  when

(a)  $Z(t) = e^{\alpha t}$ ,

(b)  $Z(t) = \int_0^t g(s)dW(s)$ , where  $g$  is an adapted stochastic process.

(c)  $Z(t) = e^{\alpha W(t)}$

(d)  $Z(t) = e^{\alpha X(t)}$ , where  $X$  has the stochastic differential

$$dX(t) = \mu dt + \sigma dW(t)$$

( $\mu$  and  $\sigma$  are constants).

(e)  $Z(t) = X^2(t)$ , where  $X$  has the stochastic differential

$$dX(t) = \alpha X(t)dt + \sigma X(t)dW(t).$$

**Exercise 1.2** Compute the stochastic differential for  $Z$  when  $Z(t) = \frac{1}{X(t)}$  and  $X$  has the stochastic differential

$$dX(t) = \alpha X(t)dt + \sigma X(t)dW(t).$$

By using the definition  $Z = X^{-1}$  you can in fact express the right hand side of  $dZ$  entirely in terms of  $Z$  itself (rather than in terms of  $X$ ). Thus  $Z$  satisfies a stochastic differential equation. Which one?

**Exercise 1.3** Let  $\sigma(t)$  be a given deterministic function of time and define the process  $X$  by

$$X(t) = \int_0^t \sigma(s) dW(s). \quad (1)$$

Show that the characteristic function of  $X(t)$  (for a fixed  $t$ ) is given by

$$E \left[ e^{iuX(t)} \right] = \exp \left\{ -\frac{u^2}{2} \int_0^t \sigma^2(s) ds \right\}, \quad u \in R, \quad (2)$$

thus showing that  $X(t)$  is normally distributed with zero mean and a variance given by

$$\text{Var}[X(t)] = \int_0^t \sigma^2(s) ds.$$

**Exercise 1.4** Suppose that  $X$  has the stochastic differential

$$dX(t) = \alpha X(t) dt + \sigma(t) dW(t),$$

where  $\alpha$  is a real number whereas  $\sigma(t)$  is any stochastic process. Determine the function  $m(t) = E[X(t)]$ .

**Exercise 1.5** Suppose that the process  $X$  has a stochastic differential

$$dX(t) = \mu(t) dt + \sigma(t) dW(t),$$

and that  $\mu(t) \geq 0$  with probability one for all  $t$ . Show that this implies that  $X$  is a submartingale.

**Exercise 1.6** A function  $h(x_1, \dots, x_n)$  is said to be **harmonic** if it satisfies the condition

$$\sum_{i=1}^n \frac{\partial^2 h}{\partial x_i^2} = 0.$$

It is **subharmonic** if it satisfies the condition

$$\sum_{i=1}^n \frac{\partial^2 h}{\partial x_i^2} \geq 0.$$

Let  $W_1, \dots, W_n$  be independent standard Wiener processes, and define the process  $X$  by  $X(t) = h(W_1(t), \dots, W_n(t))$ . Show that  $X$  is a martingale (submartingale) if  $h$  is harmonic (subharmonic).

**Exercise 1.7** The object of this exercise is to give an argument for the formal identity

$$dW_1 \cdot dW_2 = 0,$$

when  $W_1$  and  $W_2$  are independent Wiener processes. Let us therefore fix a time  $t$ , and divide the interval  $[0, t]$  into equidistant points  $0 = t_0 < t_1 < \dots < t_n = t$ , where  $t_i = \frac{i}{n} \cdot t$ . We use the notation

$$\Delta W_i(t_k) = W_i(t_k) - W_i(t_{k-1}), \quad i = 1, 2.$$

Now define  $Q_n$  by

$$Q_n = \sum_{k=1}^n \Delta W_1(t_k) \cdot \Delta W_2(t_k).$$

Show that  $Q_n \rightarrow 0$  in  $L^2$ , i.e. show that

$$\begin{aligned} E[Q_n] &= 0, \\ \text{Var}[Q_n] &\rightarrow 0. \end{aligned}$$

**Exercise 1.8** Let  $X$  and  $Y$  be given as the solutions to the following system of stochastic differential equations.

$$\begin{aligned} dX &= \alpha X dt - Y dW, & X(0) &= x_0, \\ dY &= \alpha Y dt + X dW, & Y(0) &= y_0. \end{aligned}$$

Note that the initial values  $x_0, y_0$  are deterministic constants.

- (a) Prove that the process  $R$  defined by  $R(t) = X^2(t) + Y^2(t)$  is deterministic.
- (b) Compute  $E[X(t)]$ .

**Exercise 1.9** For a  $n \times n$  matrix  $A$ , the **trace** of  $A$  is defined as

$$\text{tr}(A) = \sum_{i=1}^n A_{ii}.$$

- (a) If  $B$  is  $n \times d$  and  $C$  is  $d \times n$ , then  $BC$  is  $n \times n$ . Show that

$$\text{tr}(BC) = \sum_{ij} B_{ij} C_{ji}.$$

- (b) With assumptions as above, show that

$$\text{tr}(BC) = \text{tr}(CB).$$

(c) Show that the multi dimensional Itô formula can be written

$$df = \left\{ \frac{\partial f}{\partial t} + \sum_{i=1}^n \mu_i \frac{\partial f}{\partial x_i} + \frac{1}{2} \text{tr} [\sigma^* H \sigma] \right\} dt + \sum_{i=1}^n \frac{\partial f}{\partial x_i} \sigma_i dW_i$$

where  $H$  denotes the Hessian matrix

$$H_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}.$$

**Exercise 1.10** Show that the scalar SDE

$$\begin{aligned} dX_t &= \alpha X_t dt + \sigma dW_t, \\ X_0 &= x_0, \end{aligned}$$

has the solution

$$X(t) = e^{\alpha t} \cdot x_0 + \sigma \int_0^t e^{\alpha(t-s)} dW_s, \quad (3)$$

by differentiating  $X$  as defined by eqn (3) and showing that  $X$  so defined actually satisfies the SDE.

**Hint:** Write eqn (3) as

$$X_t = Y_t + Z_t \cdot R_t,$$

where

$$\begin{aligned} Y_t &= e^{\alpha t} \cdot x_0, \\ Z_t &= e^{\alpha t} \cdot \sigma, \\ R_t &= \int_0^t e^{-\alpha s} dW_s, \end{aligned}$$

and first compute the differentials  $dZ$ ,  $dY$  and  $dR$ . Then use the multidimensional Itô formula on the function  $f(y, z, r) = y + z \cdot r$ .

**Exercise 1.11** Let  $A$  be an  $n \times n$  matrix, and define the matrix exponential  $e^A$  by the series

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!}.$$

This series can be shown to converge uniformly.

(a) Show, by taking derivatives under the summation sign, that

$$\frac{de^{At}}{dt} = Ae^{At}.$$

(b) Show that

$$e^0 = I,$$

where  $0$  denotes the zero matrix, and  $I$  denotes the identity matrix.

(c) Convince yourself that if  $A$  and  $B$  commute, i.e.  $AB = BA$ , then

$$e^{A+B} = e^A \cdot e^B = e^B \cdot e^A.$$

**Hint:** Write the series expansion in detail.

(d) Show that  $e^A$  is invertible for every  $A$ , and that in fact

$$\left[e^A\right]^{-1} = e^{-A}.$$

(e) Show that for any  $A$ ,  $t$  and  $s$

$$e^{A(t+s)} = e^{At} \cdot e^{As}$$

(f) Show that

$$\left(e^A\right)^* = e^{A^*}$$

**Exercise 1.12** Consider the  $n$ -dimensional linear SDE

$$\begin{cases} dX_t &= (AX_t + b_t) dt + \sigma_t dW_t, \\ X_0 &= x_0 \end{cases} \quad (4)$$

where  $A$  is an  $n \times n$  matrix,  $b$  is an  $R^n$ -valued deterministic function (in column vector form),  $\sigma$  is a deterministic  $n \times d$  deterministic matrix valued function, and  $W$  an  $d$ -dimensional Wiener process. Show that the solution of this equation is given by

$$X_t = e^{At} x_0 + \int_0^t e^{A(t-s)} b_s ds + \int_0^t e^{A(t-s)} \sigma_s dW_s. \quad (5)$$

**Exercise 1.13** Consider again the linear SDE (4). Show that the expected value function

$$m(t) = E[X(t)]$$

, and the covariance matrix

$$C(t) = \{Cov(X_i(t), X_j(t))\}_{i,j}$$

are given by

$$\begin{aligned} m(t) &= e^{At} x_0 + \int_0^t e^{A(t-s)} b(s) ds, \\ C(t) &= \int_0^t e^{A(t-s)} \sigma(s) \sigma^*(s) e^{A^*(t-s)} ds, \end{aligned}$$

where  $*$  denotes transpose.

**Hint:** Use the explicit solution above, and the fact that

$$C(t) = E [X_t X_t^*] - m(t)m^*(t).$$

Geometric Brownian motion (GBM) constitutes a class of processes which is closed under a number of nice operations. Here are some examples.

**Exercise 1.14** Suppose that  $X$  satisfies the SDE

$$dX_t = \alpha X_t dt + \sigma X_t dW_t.$$

Now define  $Y$  by  $Y_t = X_t^\beta$ , where  $\beta$  is a real number. Then  $Y$  is also a GBM process. Compute  $dY$  and find out which SDE  $Y$  satisfies.

**Exercise 1.15** Suppose that  $X$  satisfies the SDE

$$dX_t = \alpha X_t dt + \sigma X_t dW_t,$$

and  $Y$  satisfies

$$dY_t = \gamma Y_t dt + \delta Y_t dV_t,$$

where  $V$  is a Wiener process which is independent of  $W$ . Define  $Z$  by  $Z = \frac{X}{Y}$  and derive an SDE for  $Z$  by computing  $dZ$  and substituting  $Z$  for  $\frac{X}{Y}$  in the right hand side of  $dZ$ . If  $X$  is nominal income and  $Y$  describes inflation then  $Z$  describes real income.

**Exercise 1.16** Suppose that  $X$  satisfies the SDE

$$dX_t = \alpha X_t dt + \sigma X_t dW_t,$$

and  $Y$  satisfies

$$dY_t = \gamma Y_t dt + \delta Y_t dW_t.$$

Note that now both  $X$  and  $Y$  are driven by the same Wiener process  $W$ . Define  $Z$  by  $Z = \frac{X}{Y}$  and derive an SDE for  $Z$ .

**Exercise 1.17** Suppose that  $X$  satisfies the SDE

$$dX_t = \alpha X_t dt + \sigma X_t dW_t,$$

and  $Y$  satisfies

$$dY_t = \gamma Y_t dt + \delta Y_t dV_t,$$

where  $V$  is a Wiener process which is independent of  $W$ . Define  $Z$  by  $Z = X \cdot Y$  and derive an SDE for  $Z$ . If  $X$  describes the price process of, for example, IBM in US\$ and  $Y$  is the currency rate SEK/US\$ then  $Z$  describes the dynamics of the IBM stock expressed in SEK.

**Exercise 1.18** Use a stochastic representation result in order to solve the following boundary value problem in the domain  $[0, T] \times \mathbb{R}$ .

$$\begin{aligned} \frac{\partial F}{\partial t} + \mu x \frac{\partial F}{\partial x} + \frac{1}{2} \sigma^2 x^2 \frac{\partial^2 F}{\partial x^2} &= 0, \\ F(T, x) &= \ln(x^2). \end{aligned}$$

Here  $\mu$  and  $\sigma$  are assumed to be known constants.

**Exercise 1.19** Consider the following boundary value problem in the domain  $[0, T] \times \mathbb{R}$ .

$$\begin{aligned} \frac{\partial F}{\partial t} + \mu(t, x) \frac{\partial F}{\partial x} + \frac{1}{2} \sigma^2(t, x) \frac{\partial^2 F}{\partial x^2} + k(t, x) &= 0, \\ F(T, x) &= \Phi(x). \end{aligned}$$

Here  $\mu$ ,  $\sigma$ ,  $k$  and  $\Phi$  are assumed to be known functions.

Prove that this problem has the stochastic representation formula

$$F(t, x) = E_{t,x} [\Phi(X_T)] + \int_t^T E_{t,x} [k(s, X_s)] ds,$$

where as usual  $X$  has the dynamics

$$\begin{aligned} dX_s &= \mu(s, X_s) ds + \sigma(s, X_s) dW_s, \\ X_t &= x. \end{aligned}$$

**Hint:** Define  $X$  as above, assume that  $F$  actually solves the PDE and consider the process  $Z_s = F(s, X_s)$ .

**Exercise 1.20** Use the result of the previous exercise in order to solve

$$\begin{aligned} \frac{\partial F}{\partial t} + \frac{1}{2} x^2 \frac{\partial^2 F}{\partial x^2} + x &= 0, \\ F(T, x) &= \ln(x^2). \end{aligned}$$

**Exercise 1.21** Consider the following boundary value problem in the domain  $[0, T] \times \mathbb{R}$ .

$$\begin{aligned} \frac{\partial F}{\partial t} + \mu(t, x) \frac{\partial F}{\partial x} + \frac{1}{2} \sigma^2(t, x) \frac{\partial^2 F}{\partial x^2} + r(t, x) F &= 0, \\ F(T, x) &= \Phi(x). \end{aligned}$$

Here  $\mu(t, x)$ ,  $\sigma(t, x)$ ,  $r(t, x)$  and  $\Phi(x)$  are assumed to be known functions. Prove that this problem has a stochastic representation formula of the form

$$F(t, x) = E_{t,x} \left[ \Phi(X_T) e^{\int_t^T r(s, X_s) ds} \right],$$

by considering the process  $Z_s = F(s, X_s) \times \exp[\int_t^s r(u, X_u) du]$  on the time interval  $[t, T]$ .

**Exercise 1.22** Solve the boundary value problem

$$\begin{aligned}\frac{\partial F}{\partial t}(t, x, y) + \frac{1}{2}\sigma^2 \frac{\partial^2 F}{\partial x^2}(t, x, y) + \frac{1}{2}\delta^2 \frac{\partial^2 F}{\partial y^2}(t, x, y) &= 0, \\ F(T, x, y) &= xy.\end{aligned}$$

**Exercise 1.23** Go through the details in the derivation of the Kolmogorov forward equation.

**Exercise 1.24** Consider the SDE

$$dX_t = \alpha dt + \sigma dW_t,$$

where  $\alpha$  and  $\sigma$  are constants.

- (a) Compute the transition density  $p(s, y; t, x)$ , by solving the SDE.
- (b) Write down the Fokker-Planck equation for the transition density and check the equation is indeed satisfied by your answer in (a).

**Exercise 1.25** Consider the standard GBM

$$dX_t = \alpha X_t dt + \sigma X_t dW_t$$

and use the representation

$$X_t = X_s \exp \left\{ \left[ \alpha - \frac{1}{2}\sigma^2 \right] (t - s) + \sigma [W_t - W_s] \right\}$$

in order to derive the transition density  $p(s, y; t, x)$  of GBM. Check that this density satisfies the Fokker-Planck equation.