

# Probability Theory and Stochastic Processes

## Solutions

### List 1

1) a) yes b) yes c) yes

3)  $2^n$

4) yes, no

5) yes

6) a) Since  $\sigma(\mathcal{A}_2)$  is a  $\sigma$ -algebra that contains  $\mathcal{A}_2$ , it also contains  $\mathcal{A}_1$ . Moreover, as  $\sigma(\mathcal{A}_1)$  is the intersection of all  $\sigma$ -algebras containing  $\mathcal{A}_1$ , it must be inside  $\sigma(\mathcal{A}_2)$ .

b) This follows because  $\sigma(\mathcal{A})$  is already a  $\sigma$ -algebra.

c) Recall that  $\mathcal{B} = \sigma(\mathcal{I})$  where  $\mathcal{I}$  is the collection of all intervals on the form  $]a, b]$ . From

$$]a, +\infty[ = \left( \bigcup_{n \in \mathbb{N}} ]-\infty, a - \frac{1}{n}] \right)^c,$$

it follows that  $\mathcal{A} \subset \sigma(\mathcal{I})$ . Furthermore,

$$\begin{aligned} ]a, b] &= ]a, +\infty[ \cap ]b, +\infty[^c \\ &= \left( \bigcup_{n \in \mathbb{N}} \left[ a + \frac{1}{n}, +\infty \right[ \right) \cap \left( \bigcup_{n \in \mathbb{N}} \left[ b + \frac{1}{n}, +\infty \right[ \right)^c \end{aligned}$$

So,  $\mathcal{I} \subset \sigma(\mathcal{A})$ . By (a),  $\sigma(\mathcal{A}) = \mathcal{B}$ .

7) see lecture notes

8) yes, no

### List 2

1)

2)

3)

4) a)

b)

5)

6) a) yes

b)  $f(a_1) + f(a_2) + f(a_3)$

7)

8)

9) a) Let  $\psi = f - g$  so that  $\psi$  is  $\mathcal{F}$ -measurable and  $\int_B \psi d\mu = 0$  for every  $B \in \mathcal{F}$ . Consider the measurable set

$$B = \{\psi^+ > 0\} \in \mathcal{F}$$

by writing  $\psi^+ = \max\{\psi, 0\}$  which is also  $\mathcal{F}$ -measurable. Notice that  $B = \{\psi = \psi^+ > 0\}$  and  $\int_B \psi d\mu = 0$ . In addition, take

$$B_n = \left\{ \psi^+ \geq \frac{1}{n} \right\} \in \mathcal{F}$$

Hence,  $B_n \uparrow B$ . So, using the Markov inequality,

$$0 \leq \mu(B_n) \leq n \int_{B_n} \psi^+ d\mu \leq n \int_B \psi^+ d\mu = n \int_B \psi d\mu = 0.$$

Then,  $\mu(B) = \lim \mu(B_n) = 0$  which means that  $\psi^+ = 0$   $\mu$ -a.e. The same idea for  $\psi^-$  implies that  $\psi = 0$   $\mu$ -a.e.

b) Let  $\Omega = [0, 1]$ ,  $\mathcal{A} = \{\emptyset, \Omega\}$  and  $\mathcal{F} = \mathcal{B}(\Omega)$ . Take the Lebesgue measure  $m$  on  $\Omega$ . Consider  $h(x) = 1/2$  and  $f(x) = x$ . For any  $A \in \mathcal{A}$ , we have  $\int_A h dm = \int_A f dm$ . However,  $f \neq h$   $m$ -a.e.

10) a) 1

b)  $1/2$

c)  $\arctan \pi$

d) 0

e) 0

### List 3

1) a)

$$F(x) = \begin{cases} 0, & x < 2 \\ \frac{1}{3}, & 2 \leq x < 3 \\ 1, & x \geq 3 \end{cases}$$

b)

$$F(x) = \begin{cases} 0, & x < 0 \\ \frac{1}{3}x, & 0 \leq x < 1 \\ \frac{1}{3}, & 1 \leq x < 2 \\ \frac{2}{3}, & 2 \leq x < 3 \\ 1, & x \geq 3 \end{cases}$$

2) a) For each  $x \in D$  we have  $F(x^-) < r_x < F(x^+)$  for some choice of a rational number  $r_x \in \mathbb{Q}$  (for example, take  $r_x$  to be the number with the smallest possible number of decimal places of  $(F(x^-) + F(x^+))/2$ ).

We can define a function  $g: D \rightarrow \mathbb{Q}$  given by  $g(x) = r_x$ . Thus, for  $x_1 < x_2$  both in  $D$  we have  $g(x_1) < F(x_1^+) \leq F(x_2^-) < g(x_2)$ , meaning that  $g$  is strictly increasing. Therefore,  $g$  is a bijection between  $D$  and  $g(D) \subset \mathbb{Q}$  and  $D$  is countable.

b)  $\alpha(\{x\}) = F(x) - F(x^-) = 0$

3)

4)

5) a)  $\phi(t) = e^{ita}$

b)  $(1 - p + pe^{it})^n$

c)  $e^{\lambda(e^{it}-1)}$

d)  $p/(1 - (1 - p)e^{it})$

e)  $pn/(1 - (1 - p)e^{it})^n$

6) a)

$$\phi(t) = \frac{e^{itb} - e^{ita}}{(b - a)it}$$

b)  $1/(1 - it)$

c)  $1/(1 + t^2)$

d)  $e^{-|t|}$

e)  $e^{it\mu - \sigma^2 t^2/2}$

#### List 4

1) a) Notice that  $f(x) \in B$  is equivalent to  $x \in f^{-1}(B)$ . So, for any  $B_1, B_2 \in \mathcal{B}$ ,

$$P(f(X) \in B_1, g(Y) \in B_2) = P(X \in f^{-1}(B_1), Y \in g^{-1}(B_2)).$$

Since  $X, Y$  are independent we get that

$$P(X \in f^{-1}(B_1), Y \in g^{-1}(B_2)) = P(X \in f^{-1}(B_1)) P(Y \in g^{-1}(B_2))$$

Thus,  $f(X), g(Y)$  are also independent.

b) Since by a)  $f(X), g(Y)$  are independent, then  $E(f(X)g(Y)) = E(f(X)) E(g(Y))$ .

2) The characteristic function is  $\phi(t) = e^{-|t|}$ . So, for  $S_n/n = (X_1 + \dots + X_n)/n$  we have the characteristic function

$$\phi_n(t) = (\phi(t/n))^n = e^{-|t|}.$$

Therefore, for any  $n$ ,  $S_n/n$  has the Cauchy distribution as well, and the limit distribution is not the normal distribution.

3)  $e^{-1}$

4) 0

5) b)  $\frac{1}{2}\mathcal{X}_C + \frac{3}{2}\mathcal{X}_{C^c}$

c) 1

**List 5**4) a) Stationary distribution:  $\alpha = (\alpha_1, \alpha_2, \dots)$  where

$$\alpha_1 = \frac{1-r}{2-r}, \quad \alpha_i = \frac{1}{2^{i-1}(2-r)}, \quad i \geq 2$$

5) a)  $S = R_+$ ,  $\tau_1 = 4 = \tau_3$ ,  $\tau_2 = 2$ b)  $S = R_+$ ,  $\tau_1 = \frac{1-p}{2}$ ,  $\tau_2 = (1-p)p$ ,  $\tau_3 = \frac{p}{2}$ ,  $\tau_4 = \frac{(1-p)^2 + p^2}{2}$